

KEY STUDIES

dōTERRA Protein

- T. Colonetti et al., "Effects of whey protein supplement in the elderly submitted to resistance training: systematic review and meta-analysis," *Int. J. Food Sci. Nutr.*, vol. 68, no. 3, pp. 257–264, May 2017, doi: 10.1080/09637486.2016.1232702.
- R. W. Davies, B. P. Carson, and P. M. Jakeman, "The Effect of Whey Protein Supplementation on the Temporal Recovery of Muscle Function Following Resistance Training: A Systematic Review and Meta-Analysis," *Nutrients*, vol. 10, no. 2, Feb. 2018, doi: 10.3390/nu10020221.
- F.-C. Lam, T. M. Khan, H. Faidah, A. Haseeb, and A. H. Khan, "Effectiveness of whey protein supplements on the serum levels of amino acid, creatinine kinase and myoglobin of athletes: a systematic review and meta-analysis," *Syst. Rev.*, vol. 8, no. 1, p. 130, 31 2019, doi: 10.1186/s13643-019-1039-z.
- K. D. Tipton, T. A. Elliott, M. G. Cree, S. E. Wolf, A. P. Sanford, and R. R. Wolfe, "Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise," *Med. Sci. Sports Exerc.*, vol. 36, no. 12, pp. 2073–2081, Dec. 2004, doi: 10.1249/01.mss.0000147582.99810.c5.
- R. W. Morton et al., "A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults," *Br. J. Sports Med.*, vol. 52, no. 6, pp. 376–384, Mar. 2018, doi: 10.1136/bjsports-2017-097608.
- F. Naclerio and E. Larumbe-Zabala, "Effects of Whey Protein Alone or as Part of a Multi-ingredient Formulation on Strength, Fat-Free Mass, or Lean Body Mass in Resistance-Trained Individuals: A Meta-analysis," *Sports Med. Auckl. NZ*, vol. 46, no. 1, pp. 125–137, Jan. 2016, doi: 10.1007/s40279-015-0403-y.
- L. H. A Castro et al., "Comparative Meta-Analysis of the Effect of Concentrated, Hydrolyzed, and Isolated Whey Protein Supplementation on Body Composition of Physical Activity Practitioners," *Nutrients*, vol. 11, no. 9, Sep. 2019, doi: 10.3390/nu11092047.
- R. E. Bergia, J. L. Hudson, and W. W. Campbell, "Effect of whey protein supplementation on body composition changes in women: a systematic review and meta-analysis," *Nutr. Rev.*, vol. 76, no. 7, pp. 539–551, 01 2018, doi: 10.1093/nutrit/nuy017.
- M. Messina, H. Lynch, J. M. Dickinson, and K. E. Reed, "No Difference Between the Effects of Supplementing With Soy Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise," *Int. J. Sport Nutr. Exerc. Metab.*, vol. 28, no. 6, pp. 674–685, Nov. 2018, doi: 10.1123/ijsnem.2018-0071.
- M. Mollahosseini, S. Shab-Bidar, M. H. Rahimi, and K. Djafarian, "Effect of whey protein supplementation on long and short term appetite: A meta-analysis of randomized controlled trials," *Clin. Nutr. ESPEN*, vol. 20, pp. 34–40, Aug. 2017, doi: 10.1016/j.clnesp.2017.04.002.
- C. H. Emilien, Y. Zhu, W. H. Hsu, P. Williamson, and J. H. Hollis, "The effect of soluble fiber dextrin on postprandial appetite and subsequent food intake in healthy adults," *Nutrition*, vol. 47, pp. 6–12, Mar. 2018, doi: 10.1016/j.nut.2017.08.016.
- S. Sindayikengera and W. Xia, "Nutritional evaluation of caseins and whey proteins and their hydrolysates from Protamex," *J. Zhejiang Univ. Sci. B*, vol. 7, no. 2, pp. 90–98, Feb. 2006, doi: 10.1631/jzus.2006. B0090.
- "Casein - an overview | ScienceDirect Topics." <https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/casein> (accessed Oct. 15, 2020).
- S. A. Jakeman et al., "Soluble corn fiber increases bone calcium retention in postmenopausal women in a dose-dependent manner: a randomized crossover trial," *Am. J. Clin. Nutr.*, vol. 104, no. 3, pp. 837–843, Sep. 2016, doi: 10.3945/ajcn.116.132761.
- C. M. Whisner et al., "Soluble maize fibre affects short-term calcium absorption in adolescent boys and girls: a randomised controlled trial using dual stable isotopic tracers," *Br. J. Nutr.*, vol. 112, no. 3, pp. 446–456, Aug. 2014, doi: 10.1017/S0007114514000981.
- C. M. Whisner et al., "Soluble Corn Fiber Increases Calcium Absorption Associated with Shifts in the Gut Microbiome: A Randomized Dose-Response Trial in Free-Living Pubertal Females," *J. Nutr.*, vol. 146, no. 7, pp. 1298–1306, 2016, doi: 10.3945/jn.115.227256.
- C. for F. S. and A. Nutrition, "Questions and Answers on Dietary Fiber," FDA, Feb. 2020, Accessed: Jul. 28, 2020. [Online]. Available: <https://www.fda.gov/food-food-labeling-nutrition/questions-and-answers-dietary-fiber>.
- R. Valcheva, N. Hotte, P. Gillevet, M. Sikaroodi, A. Thiessen, and K. L. Madsen, "Soluble Dextrin Fibers Alter the Intestinal Microbiota and Reduce Proinflammatory Cytokine Secretion in Male IL-10-Deficient Mice," *J. Nutr.*, vol. 145, no. 9, pp. 2060–2066, Sep. 2015, doi: 10.3945/jn.114.207738.
- F. Hu, Y. Niu, X. Xu, Q. Hu, Q. Su, and H. Zhang, "Resistant dextrin improves high-fat-high-fructose diet induced insulin resistance," *Nutr. Metab.*, vol. 17, May 2020, doi: 10.1186/s12986-020-00450-2.
- M. Badely, M. Sepandi, M. Samadi, K. Parastouei, and

- M. Taghdir, "The effect of whey protein on the components of metabolic syndrome in overweight and obese individuals: a systematic review and meta-analysis," *Diabetes Metab. Syndr.*, vol. 13, no. 6, pp. 3121–3131, Dec. 2019, doi: 10.1016/j.dsx.2019.11.001.
- S. Ursoniu, A. Sahebkar, F. Andrica, C. Serban, and M. Banach, "Effects of flaxseed supplements on blood pressure: A systematic review and meta-analysis of controlled clinical trial," *Clin. Nutr.*, vol. 35, no. 3, pp. 615–625, Jun. 2016, doi: 10.1016/j.clnu.2015.05.012.
- M. L. Stewart, S. D. Nikhanj, D. A. Timm, W. Thomas, and J. L. Slavin, "Evaluation of the effect of four fibers on laxation, gastrointestinal tolerance and serum markers in healthy humans," *Ann. Nutr. Metab.*, vol. 56, no. 2, pp. 91–98, 2010, doi: 10.1159/000275962.
- "CFR - Code of Federal Regulations Title 21." <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?fr=101.9&SearchTerm=nutrition%20label> (accessed Oct. 15, 2020).
- L.-M. Zhou, J.-Y. Xu, C.-P. Rao, S. Han, Z. Wan, and L.-Q. Qin, "Effect of whey supplementation on circulating C-reactive protein: a meta-analysis of randomized controlled trials," *Nutrients*, vol. 7, no. 2, pp. 1131–1143, Feb. 2015, doi: 10.3390/nu7021131.
- F. Mariotti and C. D. Gardner, "Dietary Protein and Amino Acids in Vegetarian Diets—A Review," *Nutrients*, vol. 11, no. 11, Nov. 2019, doi: 10.3390/nu1112661.
- Z. X. Lu, J. F. He, Y. C. Zhang, and D. J. Bing, "Composition, physicochemical properties of pea protein and its application in functional foods," *Crit. Rev. Food Sci. Nutr.*, vol. 60, no. 15, pp. 2593–2605, Aug. 2020, doi: 10.1080/10408398.2019.1651248.
- S. Wang, F. Zhu, and Y. Kakuda, "Sacha inchi (*Plukenetia volubilis* L.): Nutritional composition, biological activity, and uses," *Food Chem.*, vol. 265, pp. 316–328, Nov. 2018, doi: 10.1016/j.foodchem.2018.05.055.
- V. Weinborn et al., "The Effect of Plant Proteins Derived from Cereals and Legumes on Heme Iron Absorption," *Nutrients*, vol. 7, no. 11, pp. 8977–8986, Oct. 2015, doi: 10.3390/nu7115446.
- M. Lonnie, I. Laurie, M. Myers, G. Horgan, W. R. Russell, and A. M. Johnstone, "Exploring Health-Promoting Attributes of Plant Proteins as a Functional Ingredient for the Food Sector: A Systematic Review of Human Interventional Studies," *Nutrients*, vol. 12, no. 8, Jul. 2020, doi: 10.3390/nu12082291.
- A. Banaszek, J. R. Townsend, D. Bender, W. C. Vantrease, A. C. Marshall, and K. D. Johnson, "The Effects of Whey vs. Pea Protein on Physical Adaptations Following 8-Weeks of High-Intensity Functional Training (HIFT): A Pilot Study," *Sports Basel Switz.*, vol. 7, no. 1, Jan. 2019, doi: 10.3390/sports7010012.
- N. Babault et al., "Pea proteins oral supplementation promotes muscle thickness gains during resistance training: a double-blind, randomized, Placebo-controlled clinical trial vs. Whey protein," *J. Int. Soc. Sports Nutr.*, vol. 12, no. 1, p. 3, 2015, doi: 10.1186/s12970-014-0064-5.
- D. C. Nieman et al., "Effects of Whey and Pea Protein Supplementation on Post-Eccentric Exercise Muscle Damage: A Randomized Trial," *Nutrients*, vol. 12, no. 8, Aug. 2020, doi: 10.3390/nu12082382.
- J. I. Baum, B. L. Gaines, G. C. Kubas, C. F. Mitchell, and S. L. Russell, "Educational nutrition messaging at breakfast reduces snack intake and influences snack preferences in adult men and women," *Appetite*, vol. 117, pp. 67–73, 01 2017, doi: 10.1016/j.appet.2017.06.009.
- R. Abou-Samra, L. Keersmaekers, D. Brienza, R. Mukherjee, and K. Macé, "Effect of different protein sources on satiation and short-term satiety when consumed as a starter," *Nutr. J.*, vol. 10, p. 139, Dec. 2011, doi: 10.1186/1475-2891-10-139.
- S. Ibrügger, M. Kristensen, M. S. Mikkelsen, and A. Astrup, "Flaxseed dietary fiber supplements for suppression of appetite and food intake," *Appetite*, vol. 58, no. 2, pp. 490–495, Apr. 2012, doi: 10.1016/j.appet.2011.12.024.
- M. S. Abellán Ruiz et al., "[Effect of quinua (*Chenopodium quinoa*)consumption as a coadjvant in nutritional intervention in prediabetic subjects]," *Nutr. Hosp.*, vol. 34, no. 5, pp. 1163–1169, Oct. 2017, doi: 10.20960/nh.843.
- L. Li, G. Lietz, W. Bal, A. Watson, B. Morfey, and C. Seal, "Effects of Quinoa (*Chenopodium quinoa* Willd.) Consumption on Markers of CVD Risk," *Nutrients*, vol. 10, no. 6, Jun. 2018, doi: 10.3390/nu10060777.
- F. G. De Carvalho, P. P. Ovídio, G. J. Padovan, A. A. Jordão Junior, J. S. Marchini, and A. M. Navarro, "Metabolic parameters of postmenopausal women after quinoa or corn flakes intake—a prospective and double-blind study," *Int. J. Food Sci. Nutr.*, vol. 65, no. 3, pp. 380–385, May 2014, doi: 10.3109/09637486.2013.866637.
- I. Tzoulaki, P. Elliott, V. Kontis, and M. Ezzati, "Worldwide Exposures to Cardiovascular Risk Factors and Associated Health Effects: Current Knowledge and Data Gaps," *Circulation*, vol. 133, no. 23, pp. 2314–2333, Jun. 2016, doi: 10.1161/CIRCULATIONAHA.115.008718.
- K. Mosikanon, D. Arthan, A. Kettawan, R. Tungtrongchitr, and P. Prangthip, "Yeast β-Glucan Modulates Inflammation and Waist Circumference in Overweight and Obese Subjects," *J. Diet. Suppl.*, vol. 14, no. 2, pp. 173–185, Mar. 2017, doi: 10.1080/19390211.2016.1207005.
- S.-Y. Tan, P. C. Siow, E. Peh, and C. J. Henry, "Influence of rice, pea and oat proteins in attenuating glycemic response of sugar-sweetened beverages," *Eur. J.*

- Nutr., vol. 57, no. 8, pp. 2795–2803, Dec. 2018, doi: 10.1007/s00394-017-1547-3.
- B. K. Knapp, C. M. Parsons, L. L. Bauer, K. S. Swanson, and G. C. Fahey, “Soluble fiber dextrins and pullulans vary in extent of hydrolytic digestion in vitro and in energy value and attenuate glycemic and insulinemic responses in dogs,” J. Agric. Food Chem., vol. 58, no. 21, pp. 11355–11363, Nov. 2010, doi: 10.1021/jf102397r.
- R. Valcheva, N. Hotte, P. Gillevet, M. Sikaroodi, A. Thiessen, and K. L. Madsen, “Soluble Dextrin Fibers Alter the Intestinal Microbiota and Reduce Proinflammatory Cytokine Secretion in Male IL-10-Deficient Mice,” J. Nutr., vol. 145, no. 9, pp. 2060–2066, Sep. 2015, doi: 10.3945/jn.114.207738.
- M. L. Stewart, S. D. Nikhanj, D. A. Timm, W. Thomas, and J. L. Slavin, “Evaluation of the effect of four fibers on laxation, gastrointestinal tolerance and serum markers in healthy humans,” Ann. Nutr. Metab., vol. 56, no. 2, pp. 91–98, 2010, doi: 10.1159/000275962.
- K. Turunen et al., “Impact of beta-glucan on the faecal microbiota of polypectomized patients: a pilot study,” Anaerobe, vol. 17, no. 6, pp. 403–406, Dec. 2011, doi: 10.1016/j.anaerobe.2011.03.025.
- R. A. Mohamed, R. S. Ramadan, and L. A. Ahmed, “Effect of Substituting Pumpkin Seed Protein Isolate for Casein on Serum Liver Enzymes, Lipid Profile and Antioxidant Enzymes in CCl₄-intoxicated Rats,” p. 7, 2009.
- T. Dharnono, K. Rudnicka, M. Wilhelm, and C. Schoen, “Effects of Yeast (1,3)-(1,6)-Beta-Glucan on Severity of Upper Respiratory Tract Infections: A Double-Blind, Randomized, Placebo-Controlled Study in Healthy Subjects,” J. Am. Coll. Nutr., vol. 38, no. 1, pp. 40–50, 2019, doi: 10.1080/07315724.2018.1478339.
- A. Auinger, L. Riede, G. Bothe, R. Busch, and J. Gruenwald, “Yeast (1,3)-(1,6)-beta-glucan helps to maintain the body’s defence against pathogens: a double-blind, randomized, placebo-controlled, multicentric study in healthy subjects,” Eur. J. Nutr., vol. 52, no. 8, pp. 1913–1918, Dec. 2013, doi: 10.1007/s00394-013-0492-z.
- S. H. Albeituni et al., “Yeast-Derived Particulate β-Glucan Treatment Subverts the Suppression of Myeloid-Derived Suppressor Cells (MDSC) by Inducing Polymorphonuclear MDSC Apoptosis and Monocytic MDSC Differentiation to APC in Cancer,” J. Immunol. Baltim. Md 1950, vol. 196, no. 5, pp. 2167–2180, Mar. 2016, doi: 10.4049/jimmunol.1501853.
- C. Z. Nkosi, A. R. Opoku, and S. E. Terblanche, “Antioxidative effects of pumpkin seed (Cucurbita pepo) protein isolate in CCl₄-induced liver injury in low-protein fed rats,” Phytother. Res. PTR, vol. 20, no. 11, pp. 935–940, Nov. 2006, doi: 10.1002/ptr.1977.
- C. Z. Nkosi, A. R. Opoku, and S. E. Terblanche, “In Vitro antioxidative activity of pumpkin seed (Cucurbita pepo) protein isolate and its In Vivo effect on alanine transaminase and aspartate transaminase in acetaminophen-induced liver injury in low protein fed rats,” Phytother. Res. PTR, vol. 20, no. 9, pp. 780–783, Sep. 2006, doi: 10.1002/ptr.1958.
- C. for F. S. and A. Nutrition, “Review of the Scientific Evidence on the Physiological Effects of Certain Non-Digestible Carbohydrates,” FDA, Feb. 2020, Accessed: Oct. 15, 2020. [Online]. Available: <https://www.fda.gov/food/food-labeling-nutrition/review-scientific-evidence-physiological-effects-certain-non-digestible-carbohydrates>.
- C. for F. S. and A. Nutrition, “Guidance for Industry: The Declaration of Certain Isolated or Synthetic Non-Digestible Carbohydrates as Dietary Fiber on Nutrition and Supplement Facts Labels,” U.S. Food and Drug Administration, Feb. 01, 2020. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/guidance-industry-declaration-certain-isolated-or-synthetic-non-digestible-carbohydrates-dietary> (accessed Oct. 15, 2020).
- M. Kristensen et al., “Flaxseed dietary fibers lower cholesterol and increase fecal fat excretion, but magnitude of effect depend on food type,” Nutr. Metab., vol. 9, p. 8, Feb. 2012, doi: 10.1186/1743-7075-9-8.
- E. R. Grela, S. Sobolewska, and T. Roziński, “Effect of inulin extracts or inulin-containing plant supplement on blood lipid indices and fatty acid profile in fattener tissues,” Pol. J. Vet. Sci., vol. 17, no. 1, pp. 93–98, 2014, doi: 10.2478/pjvs-2014-0012.
- S. T. Auti and Y. A. Kulkarni, “Neuroprotective Effect of Cardamom Oil Against Aluminum Induced Neurotoxicity in Rats,” Front. Neurol., vol. 10, p. 399, 2019, doi: 10.3389/fneur.2019.00399.
- Due, A., Toubro, S., Skov, A. R., & Astrup, A. (2004). Effect of normal-fat diets, either medium or high in protein, on body weight in overweight subjects: a randomised 1-year trial. International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity, 28(10), 1283–1290. doi:10.1038/sj.ijo.0802767
- Epel, E., Lapidus, R., McEwen, B., & Brownell, K. (2001). Stress may add bite to appetite in women: a laboratory study of stress-induced cortisol and eating behavior. Psychoneuroendocrinology, 26(1), 37–49.
- Harding, J. L., Backholer, K., Williams, E. D., Peeters, A., Cameron, A. J., Hare, M. J., ... Magliano, D. J. (2014). Psychosocial stress is positively associated with body mass index gain over 5 years: evidence from the longitudinal AusDiab study. Obesity (Silver Spring, Md.), 22(1), 277–286. doi:10.1002/oby.20423
- Hill, A. J., Peikin, S. R., Ryan, C. A., & Blundell, J. E. (1990). Oral administration of proteinase inhibitor II from potatoes reduces energy intake in man. Physiology & Behavior, 48(2), 241–246.
- Komarnytsky, S., Cook, A., & Raskin, I. (2011). Potato protease inhibitors inhibit food intake and increase

circulating cholecystokinin levels by a trypsin-dependent mechanism. *International Journal of Obesity* (2005), 35(2), 236–243. doi:10.1038/ijo.2010.192

Newman, E., O'Connor, D. B., & Conner, M. (2007). Daily hassles and eating behaviour: the role of cortisol reactivity status. *Psychoneuroendocrinology*, 32(2), 125–132. doi:10.1016/j.psyneuen.2006.11.006

Noakes, M., Foster, P. R., Keogh, J. B., & Clifton, P. M. (2004). Meal replacements are as effective as structured weight-loss diets for treating obesity in adults with features of metabolic syndrome. *The Journal of Nutrition*, 134(8), 1894–1899.

Noakes, M., Keogh, J. B., Foster, P. R., & Clifton, P. M. (2005). Effect of an energy-restricted, high-protein, low-fat diet relative to a conventional high-carbohydrate, low-fat diet on weight loss, body composition, nutritional status, and markers of cardiovascular health in obese women. *The American Journal of Clinical Nutrition*, 81(6), 1298–1306.

Rosmond, R., Dallman, M. F., & Björntorp, P. (1998). Stress-related cortisol secretion in men: relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities. *The Journal of Clinical Endocrinology and Metabolism*, 83(6), 1853–1859. doi:10.1210/jcem.83.6.4843

Rothacker, D. Q., Staniszewski, B. A., & Ellis, P. K. (2001). Liquid meal replacement vs traditional food: a potential model for women who cannot maintain eating habit change. *Journal of the American Dietetic Association*, 101(3), 345–347. doi:10.1016/S0002-8223(01)00089-X

Schwartz, J. G., Guan, D., Green, G. M., & Phillips, W. T. (1994). Treatment with an oral proteinase inhibitor slows gastric emptying and acutely reduces glucose and insulin levels after a liquid meal in type II diabetic patients. *Diabetes Care*, 17(4), 255–262.